

# Hybrid revascularisation in multivessel coronary artery disease: could a combination of CABG and PCI be the best option in selected patients?

Benjamin J. Wrigley<sup>1\*</sup>, MBChB, MRCP; Gopal Dubey<sup>1</sup>, BSc, MBBS, MRCP; Tom Spyt<sup>2</sup>, MD, FRCS; Anthony H. Gershlick<sup>1</sup>, BSc, MBBS, FRCP

1. Department of Cardiology, University Hospitals of Leicester, Leicester, United Kingdom; 2. Department of Cardiac Surgery, University Hospitals of Leicester, Leicester, United Kingdom

## Introduction

The prognostic benefits of coronary artery bypass grafting (CABG) arise primarily from the placement of a left internal mammary artery (LIMA) onto the left anterior descending (LAD) coronary artery. In contrast, the long-term patency rates of adjunctive conduits (including vein and radial grafts) placed onto the remaining vessels are less good, with approximately half becoming occluded within 10 years. Continuing improvements in stent technology have made it possible to treat increasingly complex disease with percutaneous coronary intervention (PCI) and the long-term outcome rates with new-generation drug-eluting stents (DES) have improved significantly. Rather than considering each revascularisation technique in isolation, a hybrid approach utilises the potential benefits of both revascularisation techniques, and thus may improve long-term outcomes in selected patients. In this article, we will discuss the contemporary revascularisation strategies used in multivessel disease, review the available data for hybrid revascularisation and consider the need for a clinical trial that will provide the data most relevant to current practice.

Editorial, see page 1231

## Background

CABG is commonly reserved for patients with complex multivessel disease, especially involving the left main stem (LMS) or proximal LAD<sup>1</sup>. The excellent long-term patency rates for LIMA

grafts to the LAD are undisputed (over 95% at 10 years) and thus make this conduit the gold standard as part of a surgical revascularisation strategy<sup>2,3</sup>. Unfortunately, the long-term patency of any necessary adjunctive saphenous vein grafts (SVG) placed on the remaining coronary arteries is not so good, with only 56% of SVGs to the right coronary artery (RCA) and 58% to the circumflex artery (Cx) remaining open at 10 years<sup>4</sup>. Total arterial grafting has been increasingly used, although only 12% of all CABG procedures in the run-in phase of the SYNTAX trial used total arterial grafting<sup>5</sup>. Furthermore, the choice of arterial conduit to the non-LAD vessel may be critical, with conflicting reports on the performance of radial artery grafts versus saphenous vein grafts. There are some data to suggest lower rates of graft occlusion with radial grafts<sup>6</sup>, although more recent randomised data have shown no differences in one-year patency between radial arterial conduits and SVGs (89% for both)<sup>7</sup>. The use of bilateral internal mammary arteries (BIMA) has also been investigated, with observational studies suggesting improved mortality of BIMA compared to single LIMA grafting<sup>8</sup>. However, the widespread use of BIMA grafting has been limited by a perceived increase in procedural complexity and early morbidity, in particular sternal dehiscence, most marked in diabetic patients<sup>9,10</sup>. Consequently, if undertaken, the commonest surgical procedure for multivessel disease involves the placement of a combination of venous and

\*Corresponding author: University Hospitals of Leicester, Glenfield Hospital, Groby Road, Leicester, LE3 9QP, United Kingdom. E-mail: benwrigley@hotmail.com

arterial grafts, with 81% of surgical patients in the run-in phase of the SYNTAX study treated by this approach<sup>5</sup>.

Didactically defined improvements in PCI have ensured that a less invasive approach may be appropriate in patients with multi-vessel disease, with the emergence of DES resulting in fewer restenoses and repeat revascularisation procedures compared to balloon angioplasty and bare metal stents (BMS)<sup>11</sup>. For straightforward lesions, the target lesion failure rate (mostly restenosis requiring a repeat PCI) at one year is as low as 4-5%<sup>12</sup>.

Recognising the relative advantages and drawbacks of both CABG and PCI has led to the concept of hybrid revascularisation, which could combine the advantages of LIMA grafting to the LAD with DES-PCI to any remaining coronary vessels.

### Drug-eluting stents versus coronary artery bypass grafting

Several observational studies have compared CABG with DES placement in patients with multivessel disease. A recent pooled analysis of these studies included over 17,000 patients and showed that the risk of death with DES was similar to the risk with CABG (5.6% versus 5.9%, respectively; RR 1.18 [95% CI: 0.80-1.75];  $p=0.39$ )<sup>13</sup>. Overall MACCE rates (defined as death, myocardial infarction [MI], stroke and repeat revascularisation) were higher in DES compared to CABG (7% versus 5%, respectively,  $p=0.002$ ) being driven by higher rates of target vessel revascularisation (TVR) and higher rates of MI in PCI patients. Limitations of such observational studies, where patient characteristics often differed between the two treatment arms, suggest there is always selection bias.

The SYNTAX study was the largest prospective study with 1,800 patients with left main coronary disease and/or three-vessel disease randomised to either CABG or PCI using the TAXUS<sup>®</sup> paclitaxel-eluting stent (Boston Scientific, Natick, MA, USA)<sup>14</sup>. At one year, PCI failed to meet the prespecified margin of non-inferiority compared to CABG. The increased primary endpoint of MACCE (combined death, MI, stroke and repeat revascularisation) was higher in patients treated with PCI (17.8% versus 12.4% for CABG;  $p=0.002$ ), again driven by the need for repeat procedures in this group. Conversely, the stroke rate was significantly higher in the CABG-treated patients.

The three and five-year results of this landmark trial have recently been published and continue to demonstrate higher rates of MACCE in patients treated with PCI compared to CABG<sup>15,16</sup>. Again, one of the major driving factors was the increased need for repeat revascularisation in PCI patients compared to CABG, but there was also an increased risk of mortality and MI related to SYNTAX score, whilst stroke rates no longer differed between the two groups.

The SYNTAX score has been developed to risk stratify patients according to the severity of the coronary artery disease and this has given rise to three groups, the so-called “low-risk” patients (SYNTAX score 0-22), “intermediate-risk” patients (23-32) and “high-risk” patients ( $\geq 33$ )<sup>14</sup>. Three-year follow-up has shown no differences between CABG and PCI in patients with low SYNTAX scores but, in those with intermediate scores, MACCE is higher in PCI (27.4% versus

18.9%,  $p=0.02$ ), driven by repeat revascularisation and increased MI. For the high-risk patients, except for stroke, the MACCE and all its components (including mortality) were significantly higher in PCI compared to CABG (34.1% versus 19.5%,  $p<0.001$ ).

It would appear that CABG is the dominant revascularisation strategy for those with high ( $\geq 33$ ) SYNTAX scores, whereas those with low scores are likely to undergo PCI due to comparable outcomes with surgery. For patients with intermediate SYNTAX scores, CABG may have an early advantage over PCI with lower rates of revascularisation at three-year follow-up. However, the majority (81% in the SYNTAX trial run-in phase<sup>5</sup>) of surgical patients receive a combination of arterial and venous grafts. Therefore, it is too soon to say whether long-term revascularisation outcomes (e.g., at 10 years) will indeed ultimately favour CABG. Further, the DES used was one that has repeatedly been shown to be inferior, especially in terms of the need for repeat revascularisation compared to the currently used new-generation DES<sup>12</sup>.

### Drug-eluting stents compared with non-left internal mammary artery conduits

As previously discussed, a major weakness of CABG revascularisation is the attrition of SVGs<sup>17-19</sup>. In spite of only 10.7% of CABG patients requiring repeat revascularisation at three years in the SYNTAX study<sup>14</sup>, this figure is likely to increase over time as vein grafts deteriorate. At ten years, published data indicate that this figure could be as high as 50%<sup>4</sup>, although the contemporary use of aggressive statin therapy following CABG may improve long-term SVG patency<sup>20</sup>. DES have been in use for ten years, although there have been quantum changes in polymer bioneutrality, stent design and drug choice between old and new-generation stents which make long-term analysis of results less meaningful than the unchanged LIMA to LAD procedure. In the RAVEL study, the first-generation Cypher<sup>®</sup> sirolimus-eluting stent (Cordis, Johnson & Johnson, Warren, NJ, USA) was associated with 89.7% freedom from target lesion revascularisation (TLR) at five years<sup>21</sup> and, in the j-Cypher registry of almost 20,000 lesions, TLR at five years was 15.9%<sup>22</sup>. The incidence within the first year was 7.3%, followed by an additional 2.2% per year for the remaining four years of follow-up. Ten-year outcome data from the DESIRE registry of 4,000 patients receiving DES since 2002 have shown target vessel revascularisation rates of only 5.3%<sup>23</sup>. While one might speculate that in the longer term contemporary DES may well be superior to the long-term results of SVGs, it must be emphasised that there have been no head-to-head randomised comparisons between DES and SVGs. With new-generation stents, the incidences of restenosis and stent thrombosis are further reduced, in part due to the cobalt-chromium platform enabling thinner stent struts. The rates of clinical restenosis at two years in the large observational Swedish Coronary Angiography and Angioplasty Register (SCAAR) were 3.9% in new-generation DES compared to 5.8% in old-generation DES (and 7.4% in bare metal stents)<sup>24</sup>. Four-year data for the XIENCE everolimus-eluting stent (Abbott Vascular, Santa Clara, CA, USA) are available from the SPIRIT II

trial, which randomised patients to XIENCE or TAXUS stents<sup>25</sup>. There was a trend towards lower rates of TLR with XIENCE (5.9% versus 12.7%,  $p=0.07$ ), with very low rates (0.9%) of definite or probable stent thrombosis.

Thus, it might be reasonably argued that, with the contemporary longevity of DES but with no advances in conduit use, PCI would be superior to SVGs over the longer term. Coupled with the excellent patency of LIMA-LAD grafts, this has led to the under-considered concept of a hybrid approach to revascularisation.

## Hybrid revascularisation

To date, the surgical aspect of the hybrid approach has typically used an “off-pump” strategy, which in some studies has been shown to reduce the risk of complications, including stroke and neurocognitive impairment<sup>26,27</sup>. Moreover, due to its anatomical course, the LIMA can be mobilised using minimally invasive direct (MID) techniques which appear under a number of guises, including MID-CAB, thoracoscopic MIDCAB, robotic MIDCAB and totally endoscopic MIDCAB<sup>28</sup>. Experienced operators have been able to achieve equivalent long-term outcomes compared to conventional LIMA-LAD grafting<sup>29-31</sup>.

There have been numerous reports of successful hybrid revascularisation in patients with multivessel disease<sup>32-35</sup>, although only five small studies to date have compared the hybrid approach to traditional CABG and none of these was a randomised trial (Table 1). The

first so-called “comparative” study was performed by de Canniere et al in 2001, where 20 patients with two-vessel coronary disease underwent hybrid revascularisation (CABG+PCI), with an interval between LIMA-LAD grafting and PCI of one to three days<sup>36</sup>. Comparisons were made with a group of 20 matched patients (according to age, sex, comorbidities, coronary anatomy and ejection fraction) undergoing conventional CABG. The hybrid procedure group had an event-free rate of 85%, compared to 35% in CABG arm (with defined events including episodes of atrial fibrillation, pericardial effusion, MI, blood transfusion requirement and leg wound dehiscence). Recovery was also more rapid in the hybrid group, with patients returning to work after  $22\pm 8$  days compared to  $89\pm 22$  days in conventional CABG patients ( $p<0.005$ ). In 2008, Kon et al performed a study using a truly simultaneous approach, whereby MIDCAB LIMA-LAD grafting was immediately followed by PCI (with DES) to the remaining vessels<sup>35</sup>. All hybrid patients ( $n=15$ ) received aspirin (325 mg) before the procedure, heparin during surgery (with no reversal on completion) and clopidogrel loading (300 mg) via a nasogastric tube on returning to the intensive care unit. They compared the hybrid procedures with 30 parallel matched controls and found that the hybrid approach resulted in no postoperative MACE (defined as MI, stroke, death or repeat intervention) compared to seven events in the OPCAB group (six MI, one stroke,  $p=0.05$ ). Long-term graft patency was assessed using CT angiography and demonstrated one stent failure in the

**Table 1. Non-randomised studies comparing hybrid revascularisation with conventional coronary artery bypass grafting.**

Study	Timing of hybrid	Method of hybrid surgery	Method of surgery in controls	No. of hybrid	No. of controls	Results
Halkos 2011	Staged	MIDCAB	OPCAB	147	588	Median follow-up 3.2 years. Primary endpoint MACCE (death, MI, stroke) similar between the groups. Repeat revascularisation higher in hybrid (12.2% vs. 3.7%, $p<0.001$ ) Similar estimated 5-year survival between groups
Vassiliades 2009	Staged	Endo-CAB	OPCAB	91	4,175	Follow-up 12 months. Primary endpoint Similar 30-day MACCE and mortality rates Total MACCE (death/stroke MI/TVR) at 12 months 10% LIMA patency 100%
Kon 2008	Simultaneous	MIDCAB	OPCAB	15	30	Follow-up 12 months Primary endpoint In-hospital MACE (death, MI, stroke, repeat revascularisation) 0 (hybrid) versus 7 (control) $p=0.05$ Secondary endpoint Hospital stay= $3.7\pm 1.4$ days vs. $6.4\pm 2.2$ days, $p<0.001$ Blood transfusion= $0.2$ units vs. $1.4$ units $p<0.001$ LIMA patency 100% MACE at 12 months= $7\%$ vs. $23\%$ $p<0.05$
Reicher 2008	Simultaneous	MIDCAB	OPCAB	13	26	Primary endpoint Mortality - no deaths in either group Hospital stay= $3.6\pm 1.5$ vs. $6.3\pm 2.3$ days, $p<0.001$ Intubation time= $0.5\pm 1.3$ vs. $11.7\pm 9.6$ hrs, $p<0.02$ Blood transfusion= $0.33\pm 0.49$ vs. $1.47\pm 1.53$ , $p<0.01$ Secondary endpoint Pre-discharge LIMA patency= $100\%$
de Canniere 2001	Mixed	MIDCAB	on-pump	20	20	Follow-up 24 months LIMA patency 100% ITU stay= $20.35\pm 2.7$ vs. $28.6\pm 12.31$ hours Hospital stay= $6.7\pm 0.7$ vs. $9.0\pm 1.2$ days, $p<0.05$

hybrid group compared to seven SVG failures in the OPCAB group ( $p=0.062$ ). In a similar study in 2008, Reicher et al looked at 13 patients undergoing simultaneous hybrid revascularisation and compared them to 26 propensity score matched parallel controls undergoing off-pump CABG<sup>37</sup>. There were no deaths in either group, but hybrid patients had a shorter length of stay ( $3.6\pm 1.5$  versus  $6.3\pm 2.3$  days,  $p=0.0001$ ). Interestingly, hybrid patients required fewer blood transfusions than controls (21% versus 59%,  $p=0.05$ ) despite the more aggressive antithrombotic regimen against a background of surgery in this group.

On a larger scale, Vassiliades et al compared 91 hybrid patients with a total 4,175 OPCAB procedures, making adjustments for selection bias using propensity scoring<sup>38</sup>. In contrast to the earlier studies, simultaneous revascularisation with surgery and PCI was not performed. Instead, 85 (93.4%) patients had LIMA to LAD grafting first, followed by PCI 2.2 $\pm$ 1.2 days later. The remainder underwent PCI first, with surgery being performed 67.8 $\pm$ 39.9 days later, and patients remained on clopidogrel throughout. Thirty-day MACCE (defined as death, MI, stroke and need for repeat intervention) rates were similar between the two study groups (1.1% in hybrid versus 3.0% in OPCAB,  $p=0.48$ ). Kaplan-Meier survival estimates at three years did not differ significantly between groups ( $p=0.14$ ).

The largest comparison of hybrid surgery with conventional OPCAB was conducted by Halkos and colleagues in 2011<sup>39</sup>. A total of 147 patients underwent hybrid revascularisation (using endoscopic CAB  $\pm$  robotic assistance + PCI) and were matched with 588 patients undergoing OPCAB (4:1 ratio) using an optimal matching algorithm. Patients were only included if the cardiologists felt that a “technically excellent” result from non-LAD stenting was possible, which generally excluded long lesions requiring multiple stents, small-calibre vessels, bifurcation lesions and chronic total occlusions (CTOs). In the Halkos study, fewer than 10 patients underwent simultaneous hybrid procedures, with the majority having surgery followed by PCI two to three days later, although some patients had PCI first if the non-LAD lesions were deemed “critical” (not clearly defined in the paper) in a joint decision between surgeon and interventional cardiologist. Patients undergoing surgery after PCI did so without interruption of clopidogrel. In-hospital outcomes were similar, with a 2% MACCE rate (composite death, stroke and MI) in each group. Repeat revascularisation was



performed if clinically indicated and was higher in the hybrid group (12.2% versus 3.7%,  $p<0.001$ ). This was driven by the need to treat LIMA-LAD lesions, progression of native disease and in-stent restenoses (3.4%). Only 2.4% of SVGs required intervention. Despite this, there were no differences in the estimated five-year survival (86.8% for the hybrid group and 84.3% for OPCAB group,  $p=0.61$ ).

The results of the above trials demonstrate the feasibility of the hybrid technique, but many questions remain unanswered, including whether it is really worth considering as an approach, which patients may benefit most from this strategy, the logistics of performing two procedures instead of one, and the safety concerns regarding the choice of antiplatelet therapy.

## Using hybrid revascularisation in clinical practice

There are several patterns of disease that might be best treated with the hybrid approach but there are a number of issues which have yet to be addressed (**Table 2**), with the optimal order in which surgery and PCI should be performed being perhaps the most important consideration. In most studies to date the LAD has been LIMA-grafted first followed by PCI to the remaining vessels. This avoids performing surgery on patients who are taking clopidogrel, which is known to increase the risks of bleeding<sup>40</sup>. Of course there may be a perceived possibility of increased perioperative risk when performing incomplete surgical revascularisation in patients with critical RCA or Cx disease if surgery is to be performed before PCI or even failure of PCI to the non-LAD lesions. However, a recent large observational study suggested that the early and long-term survival was similar with incomplete compared to complete revascularisation, so long as a LIMA was placed onto the LAD<sup>41</sup>. If PCI was to be performed first, patients would either have to undergo surgery on clopidogrel, have it stopped for five to seven days or wait several months before it could be safely and permanently discontinued. A potential alternative might be to bridge with an intravenous P2Y12 receptor antagonist, such as cangrelor. This novel agent has been studied in patients who require interruption of a thienopyridine for CABG and may be a potential solution for those requiring surgery after PCI, although further data are required<sup>42</sup>. An oral version, ticagrelor, has been used in operative patients and is reversible and

**Table 2. The pros and cons of choosing either surgery or PCI as the initial revascularisation strategy as part of the hybrid approach.**

	PCI undertaken before surgery	Surgery undertaken before PCI
		
PROS	Reduced ischaemic burden during subsequent CABG – Potentially lower perioperative risk	No need to interrupt clopidogrel after PCI Possibility of distal LMS lesions to be treated with a single stent into circumflex (as the LAD is “protected”) LIMA graft patency can be assessed at time of PCI
CONS	Perioperative bleeding/stent thrombosis – Increased perioperative bleeding if clopidogrel not discontinued – Risk of stent thrombosis if clopidogrel interrupted	Failure of PCI may necessitate repeat surgery Perceived increased perioperative ischaemia from ungrafted non-LAD lesions



could be an alternative to clopidogrel in the hybrid patients. Finally, if surgery is performed first, one must be fairly confident that PCI is technically achievable, since subjecting a patient to repeat surgery for failed PCI would clearly increase the risk of morbidity and mortality.

Most of the hybrid data so far have been generated in centres with great expertise in performing minimally invasive LIMA grafting and even using hybrid operating rooms capable of facilitating combined surgery and PCI. This is clearly not consistent with global surgical and interventional practice. Furthermore, data from the most recent observational study suggest that the need for repeat revascularisation is higher when the LIMA-LAD graft is placed using a minimally invasive approach<sup>39</sup>. Consequently, there may be a reluctance among many cardiac surgeons to adopt a technique that appears to have a steep learning curve, involves longer operating times and may not achieve the same results as conventional surgery<sup>43</sup>.

There is therefore a need to develop a standard and universally accessible strategy to manage patients with RCA and Cx disease suitable for PCI and LAD disease suitable for LIMA.

### Hybrid strategy: time for a trial?

Despite randomised studies comparing CABG and DES in multivessel disease, the optimal revascularisation strategy is far from clear and there are limitations with the SYNTAX trial that make it fall short of being the definitive study. Although the need for repeat revascularisation was higher with PCI than CABG at three years, long-term SVG patency data from other studies might indicate that revascularisation with CABG is likely to be higher over the longer-term follow-up. Furthermore, there was no mandate to use ischaemia testing to guide revascularisation, whereas the FAME trial (angiographic versus fractional flow reserve [FFR]-guided PCI) showed that FFR-guided PCI resulted in fewer stents being implanted, with a 30% reduction in the risk of death, MI or repeat revascularisation at one year<sup>44</sup>. Consequently, PCI may have been performed on non-flow-limiting lesions in the SYNTAX trial, perhaps reflected by one third of patients receiving more than 100 mm of stent. However, it should be acknowledged that FFR-guided CABG may also improve outcomes from surgery, perhaps by restricting the placement of grafts onto coronary vessels with functionally significant stenoses only.

Moving beyond the concept of hybrid revascularisation utilising a minimally invasive approach, perhaps a question more pertinent to real-world practice might be “could a hybrid strategy of conventional surgery to place LIMA-LAD (including full sternotomy/on-pump) plus DES-PCI to other vessels be a better way of managing multivessel disease than standard CABG?” One small non-randomised study has examined this strategy in 18 patients undergoing conventional CABG (majority on-pump) to place LIMA-LAD followed by DES-PCI to the remaining vessels 48 hours after surgery<sup>45</sup>. Comparisons were made with 18 matched controls (for baseline clinical characteristics and SYNTAX scores) undergoing conventional CABG with LIMA-LAD and at least one additional graft. The hybrid procedure was associated with shorter durations

of cardiopulmonary bypass and there were no differences in bleeding or duration of hospital stay. One-year MACE rates (death, MI, target vessel revascularisation) were similar.

An appropriately sized randomised trial with relevance to the majority of patients requiring revascularisation for multivessel disease is probably needed, if only to put the question to bed. The “Could Hybrid In Multivessel disease be the Rational Approach (CHIMERA)” study is being planned. Patients with multivessel disease amenable to both PCI and CABG would be randomised to either 1) conventional surgery (LIMA-LAD and either SVG/arterial conduits or RIMA to remaining vessels); 2) PCI to all vessels with contemporary DES (complemented by ischaemia testing); or 3) a hybrid approach using any strategy that places a LIMA graft onto the LAD (conventional sternotomy, on or off-pump technique or minimally invasive approach) and DES-PCI to the remaining vessels. Such a study may help patients to determine the best strategy to improve outcomes in the long term.

### Conclusions

Whilst the long-term patency of LIMA to LAD grafts is excellent, SVGs placed on the remaining vessels are disappointing and the relatively short follow-up period for patients in randomised trials does not allow for the published attrition of SVGs. New-generation stents are associated with reduced rates of restenosis and thrombosis and, although there are no randomised trials comparing DES with SVGs, it is conceivable that stents may prove superior to vein grafts over a long follow-up period. The optimal revascularisation strategy in patients with multivessel disease is therefore still not fully resolved, not least because of the variation in disease patterns that make up “multivessel disease”. A hybrid approach combines the advantages of both CABG and PCI, although minimally invasive techniques used hitherto are not reflective of current surgical practice. Studies are badly needed to evaluate whether this approach can translate into better long-term outcomes for patients with multivessel disease.

### Conflict of interest statement

A. Gershlick is on the advisory boards of, and on the lecture circuit for, Medtronic, Abbott, Boston Scientific and The Medicines Company. The other authors have no conflicts of interest to declare.

### References

1. Wijns W, Kolh P, Danchin N, Di Mario C, Falk V, Folliguet T, Garg S, Huber K, James S, Knuuti J, Lopez-Sendon J, Marco J, Menicanti L, Ostojic M, Piepoli MF, Pirlet C, Pomar JL, Reifart N, Ribichini FL, Schalij MJ, Sergeant P, Serruys PW, Silber S, Sousa Uva M, Taggart D. Guidelines on myocardial revascularization. *Eur Heart J*. 2010;31:2501-55.
2. Cameron A, Davis KB, Green G, Schaff HV. Coronary bypass surgery with internal-thoracic-artery grafts--effects on survival over a 15-year period. *N Engl J Med*. 1996;334:216-9.
3. Loop FD, Lytle BW, Cosgrove DM, Stewart RW, Goormastic M, Williams GW, Golding LA, Gill CC, Taylor PC, Sheldon WC. Influence

of the internal-mammary-artery graft on 10-year survival and other cardiac events. *N Engl J Med.* 1986;314:1-6.

4. Goldman S, Zadina K, Moritz T, Ovitt T, Sethi G, Copeland JG, Thottapurathu L, Krasnicka B, Ellis N, Anderson RJ, Henderson W. Long-term patency of saphenous vein and left internal mammary artery grafts after coronary artery bypass surgery: results from a Department of Veterans Affairs Cooperative Study. *J Am Coll Cardiol.* 2004;44:2149-56.

5. Kappetein AP, Dawkins KD, Mohr FW, Morice MC, Mack MJ, Russell ME, Pomar J, Serruys PW. Current percutaneous coronary intervention and coronary artery bypass grafting practices for three-vessel and left main coronary artery disease. Insights from the SYNTAX run-in phase. *Eur J Cardiothorac Surg.* 2006;29:486-91.

6. Desai ND, Cohen EA, Naylor CD, Fremes SE. A randomized comparison of radial-artery and saphenous-vein coronary bypass grafts. *N Engl J Med.* 2004;351:2302-9.

7. Goldman S, Sethi GK, Holman W, Thai H, McFalls E, Ward HB, Kelly RF, Rhenman B, Tobler GH, Bakaeen FG, Huh J, Soltero E, Moursi M, Haime M, Crittenden M, Kasirajan V, Ratliff M, Pett S, Irampen A, Gunnar W, Thomas D, Fremes S, Moritz T, Reda D, Harrison L, Wagner TH, Wang Y, Planting L, Miller M, Rodriguez Y, Juneman E, Morrison D, Pierce MK, Kreamer S, Shih MC, Lee K. Radial artery grafts vs saphenous vein grafts in coronary artery bypass surgery: a randomized trial. *JAMA.* 2011;305:167-74.

8. Taggart DP, D'Amico R, Altman DG. Effect of arterial revascularisation on survival: a systematic review of studies comparing bilateral and single internal mammary arteries. *Lancet.* 2001;358:870-5.

9. Ioannidis JP, Galanos O, Katritsis D, Connery CP, Drossos GE, Swistel DG, Anagnostopoulos CE. Early mortality and morbidity of bilateral versus single internal thoracic artery revascularization: propensity and risk modeling. *J Am Coll Cardiol.* 2001;37:521-8.

10. Taggart DP, Altman DG, Gray AM, Lees B, Nugara F, Yu LM, Campbell H, Flather M. Randomized trial to compare bilateral vs. single internal mammary coronary artery bypass grafting: 1-year results of the Arterial Revascularisation Trial (ART). *Eur Heart J.* 2010;31:2470-81.

11. Stettler C, Wandel S, Allemann S, Kastrati A, Morice MC, Schomig A, Pfisterer ME, Stone GW, Leon MB, de Lezo JS, Goy JJ, Park SJ, Sabate M, Suttorp MJ, Kelbaek H, Spaulding C, Menichelli M, Vermeersch P, Dirksen MT, Cervinka P, Petronio AS, Nordmann AJ, Diem P, Meier B, Zwahlen M, Reichenbach S, Trelle S, Windecker S, Juni P. Outcomes associated with drug-eluting and bare-metal stents: a collaborative network meta-analysis. *Lancet.* 2007;370:937-48.

12. Stone GW, Rizvi A, Newman W, Mastali K, Wang JC, Caputo R, Doostzadeh J, Cao S, Simonton CA, Sudhir K, Lansky AJ, Cutlip DE, Kereiakes DJ. Everolimus-eluting versus paclitaxel-eluting stents in coronary artery disease. *N Engl J Med.* 2010;362:1663-74.

13. From AM, Al Badarin FJ, Cha SS, Rihal CS. Percutaneous coronary intervention with drug-eluting stents versus coronary artery bypass surgery for multivessel coronary artery disease: a meta-analysis of data from the ARTS II, CARDia, ERACI III, and

SYNTAX studies and systematic review of observational data. *EuroIntervention.* 2010;6:269-76.

14. Serruys PW, Morice MC, Kappetein AP, Colombo A, Holmes DR, Mack MJ, Stahle E, Feldman TE, van den Brand M, Bass EJ, Van DN, Leadley K, Dawkins KD, Mohr FW. Percutaneous coronary intervention versus coronary-artery bypass grafting for severe coronary artery disease. *N Engl J Med.* 2009;360:961-72.

15. Mohr F, Redwood S, Venn G, Colombo A, Mack M, Kappetein P, Morice MC, Feldman T, Stahle E, Dawkins K, Serruys PW. TCT-43 Final five-year follow-up of the SYNTAX Trial: optimal revascularization strategy in patients with three-vessel disease. *J Am Coll Cardiol.* 2012;60 (17\_S)

16. Kappetein AP, Feldman TE, Mack MJ, Morice MC, Holmes DR, Stahle E, Dawkins KD, Mohr FW, Serruys PW, Colombo A. Comparison of coronary bypass surgery with drug-eluting stenting for the treatment of left main and/or three-vessel disease: 3-year follow-up of the SYNTAX trial. *Eur Heart J.* 2011;32:2125-34.

17. Fitzgibbon GM, Kafka HP, Leach AJ, Keon WJ, Hooper GD, Burton JR. Coronary bypass graft fate and patient outcome: angiographic follow-up of 5,065 grafts related to survival and reoperation in 1,388 patients during 25 years. *J Am Coll Cardiol.* 1996;28:616-26.

18. Gansera B, Schmidler F, Angelis I, Kiask T, Kemkes BM, Botzenhardt F. Patency of internal thoracic artery compared to vein grafts - postoperative angiographic findings in 1189 symptomatic patients in 12 years. *Thorac Cardiovasc Surg.* 2007;55:412-7.

19. Alexander JH, Hafley G, Harrington RA, Peterson ED, Ferguson TB Jr, Lorenz TJ, Goyal A, Gibson M, Mack MJ, Gennevois D, Califf RM, Kouchoukos NT. Efficacy and safety of edifoligide, an E2F transcription factor decoy, for prevention of vein graft failure following coronary artery bypass graft surgery: PREVENT IV: a randomized controlled trial. *JAMA.* 2005;294:2446-54.

20. Kulik A, Voisine P, Mathieu P, Masters RG, Mesana TG, Le May MR, Ruel M. Statin therapy and saphenous vein graft disease after coronary bypass surgery: analysis from the CASCADE randomized trial. *Ann Thorac Surg.* 2011;92:1284-90.

21. Morice MC, Serruys PW, Sousa JE, Fajadet J, Ban HE, Perin M, Colombo A, Schuler G, Barragan P, Guagliumi G, Molnar F, Falotico R. A randomized comparison of a sirolimus-eluting stent with a standard stent for coronary revascularization. *N Engl J Med.* 2002;346:1773-80.

22. Kimura T, Morimoto T, Nakagawa Y, Kawai K, Miyazaki S, Muramatsu T, Shiode N, Namura M, Sone T, Oshima S, Nishikawa H, Hiasa Y, Hayashi Y, Nobuyoshi M, Mitudo K. Very late stent thrombosis and late target lesion revascularization after sirolimus-eluting stent implantation: five-year outcome of the j-Cypher Registry. *Circulation.* 2012;125:584-91.

23. Costa JR, Sousa A, Moreira A, Costa R, Cano M, Maldonado G, Palmieri B, Sousa JE. Ten-year follow-up of the DESIRE registry: A single-center perspective on drug-eluting stents a decade later. *J Am Coll Cardiol.* 2012;59:E324.

24. Sarno G, Lagerqvist B, Frobert O, Nilsson J, Olivecrona G, Omerovic E, Saleh N, Venetznos D, James S. Lower risk of stent

thrombosis and restenosis with unrestricted use of 'new-generation' drug-eluting stents: a report from the nationwide Swedish Coronary Angiography and Angioplasty Registry (SCAAR). *Eur Heart J*. 2012;33:606-13.

25. Garg S, Serruys PW, Miquel-Hebert K. Four-year clinical follow-up of the XIENCE V everolimus-eluting coronary stent system in the treatment of patients with de novo coronary artery lesions: the SPIRIT II trial. *Catheter Cardiovasc Interv*. 2011;77:1012-7.

26. Afilalo J, Rasti M, Ohayon SM, Shimony A, Eisenberg MJ. Off-pump vs. on-pump coronary artery bypass surgery: an updated meta-analysis and meta-regression of randomized trials. *Eur Heart J*. 2012;33:1257-67.

27. Zamvar V, Williams D, Hall J, Payne N, Cann C, Young K, Karthikeyan S, Dunne J. Assessment of neurocognitive impairment after off-pump and on-pump techniques for coronary artery bypass graft surgery: prospective randomised controlled trial. *BMJ*. 2002;325:1268.

28. DeRose JJ. Current state of integrated "hybrid" coronary revascularization. *Semin Thorac Cardiovasc Surg*. 2009;21:229-36.

29. Presbitero P, Nicolini F, Maiello L, Franciosi G, Carcagni A, Milone F, Manasse E, Gallotti R. "Hybrid" percutaneous and surgical coronary revascularization: selection criteria from a single-center experience. *Ital Heart J*. 2001;2:363-8.

30. Us MH, Basaran M, Yilmaz M, Yaymaci B, Ulusoy E, Sanioglu S, Ozbek C, Arslan Y, Pohan S, Yilmaz AT. Hybrid coronary revascularization in high-risk patients. *Tex Heart Inst J*. 2006;33:458-62.

31. Repossini A, Moriggia S, Cianci V, Parodi O, Sganzerla P, Baldrighi G, Bortone F, Arena V. The LAST operation is safe and effective: MIDCABG clinical and angiographic evaluation. *Ann Thorac Surg*. 2000;70:74-8.

32. Bonatti J, Schachner T, Bonaros N, Jonetzko P, Ohlinger A, Ruetzler E, Kolbitsch C, Feuchtner G, Laufer G, Pachinger O, Friedrich G. Simultaneous hybrid coronary revascularization using totally endoscopic left internal mammary artery bypass grafting and placement of rapamycin eluting stents in the same interventional session. The COMBINATION pilot study. *Cardiology*. 2008;110:92-5.

33. Katz MR, Van Praet F, de Canniere D, Murphy D, Siwek L, Seshadri-Kreaden U, Friedrich G, Bonatti J. Integrated coronary revascularization: percutaneous coronary intervention plus robotic totally endoscopic coronary artery bypass. *Circulation*. 2006;114:1473-6.

34. Kiaii B, McClure RS, Stewart P, Rayman R, Swinamer SA, Suematsu Y, Fox S, Higgins J, Albion C, Kostuk WJ, Almond D, Sridhar K, Teffy P, Jablonsky G, Diamantouros P, Dobkowski WB, Jones P, Bainbridge D, Iglesias I, Murkin J, Cheng D, Novick RJ. Simultaneous integrated coronary artery revascularization with long-term angiographic follow-up. *J Thorac Cardiovasc Surg*. 2008;136:702-8.

35. Kon ZN, Brown EN, Tran R, Joshi A, Reicher B, Grant MC, Kallam S, Burris N, Connerney I, Zimrin D, Poston RS. Simultaneous hybrid coronary revascularization reduces postoperative morbidity

compared with results from conventional off-pump coronary artery bypass. *J Thorac Cardiovasc Surg*. 2008;135:367-75.

36. de Canniere D, Jansens JL, Goldschmidt-Clermont P, Barvais L, Decroly P, Stoupe E. Combination of minimally invasive coronary bypass and percutaneous transluminal coronary angioplasty in the treatment of double-vessel coronary disease: Two-year follow-up of a new hybrid procedure compared with "on-pump" double bypass grafting. *Am Heart J*. 2001;142:563-70.

37. Reicher B, Poston RS, Mehra MR, Joshi A, Odonkor P, Kon Z, Reyes PA, Zimrin DA. Simultaneous "hybrid" percutaneous coronary intervention and minimally invasive surgical bypass grafting: feasibility, safety, and clinical outcomes. *Am Heart J*. 2008;155:661-7.

38. Vassiliades TA, Kilgo PD, Douglas JS, Babaliaros VC, Block PC, Samady H, Cates CU, Rab ST, Morris DC. Clinical outcomes after hybrid coronary revascularisation versus off-pump coronary bypass: a prospective evaluation. *Innovations (Phila)*. 2009;4:299-306.

39. Halkos ME, Vassiliades TA, Douglas JS, Morris DC, Rab ST, Liberman HA, Samady H, Kilgo PD, Guyton RA, Puskas JD. Hybrid coronary revascularization versus off-pump coronary artery bypass grafting for the treatment of multivessel coronary artery disease. *Ann Thorac Surg*. 2011;92:1695-701.

40. Berger JS, Frye CB, Harshaw Q, Edwards FH, Steinhubl SR, Becker RC. Impact of clopidogrel in patients with acute coronary syndromes requiring coronary artery bypass surgery: a multicenter analysis. *J Am Coll Cardiol*. 2008;52:1693-701.

41. Rastan AJ, Walther T, Falk V, Kempfert J, Merk D, Lehmann S, Holzhey D, Mohr FW. Does reasonable incomplete surgical revascularization affect early or long-term survival in patients with multivessel coronary artery disease receiving left internal mammary artery bypass to left anterior descending artery? *Circulation*. 2009;120:S70-7.

42. Angiolillo DJ, Firstenberg MS, Price MJ, Tummala PE, Hutyla M, Welsby IJ, Voeltz MD, Chandna H, Ramaiah C, Brtko M, Cannon L, Dyke C, Liu T, Montalescot G, Manoukian SV, Prats J, Topol EJ. Bridging antiplatelet therapy with cangrelor in patients undergoing cardiac surgery: a randomized controlled trial. *JAMA*. 2012;307:265-74.

43. Gersbach P, Imsand C, von Segesser LK, Delabays A, Vogt P, Stumpe F. Beating heart coronary artery surgery: is sternotomy a suitable alternative to minimal invasive technique? *Eur J Cardiothorac Surg*. 2001;20:760-4.

44. Tonino PA, De BB, Pijls NH, Siebert U, Ikeno F, Veer M, Klauss V, Manoharan G, Engstrom T, Oldroyd KG, Ver Lee PN, MacCarthy PA, Fearon WF. Fractional flow reserve versus angiography for guiding percutaneous coronary intervention. *N Engl J Med*. 2009;360:213-24.

45. Delhaye C, Sudre A, Lemesle G, Vanesson L, Koussa M, Fayad G, Bauters C, Lablanche JM, Modine T. Hybrid revascularization, comprising coronary artery bypass graft with exclusive arterial conduits followed by early drug-eluting stent implantation, in multivessel coronary artery disease. *Arch Cardiovasc Dis*. 2010;103:502-11.